

NEUTRINO INTERACTIONS AT MINIBOONE

Jennifer L. Raaf
February 12, 2004
NOON 2004 Workshop



OVERVIEW

MiniBooNE

Beam and Detector

Event Reconstruction and Particle ID

Calibration and Neutrino Data

Data/MC Comparisons

Charged current quasi-elastic events

Neutral current π^0 events

Neutral current elastic events


Conclusions

MINIBOONE MOTIVATION

LSND: 3.8σ $\bar{\nu}_e$ excess from $\bar{\nu}_\mu$ source

Oscillation probability:
 $(0.264 \pm 0.067 \pm 0.045)\%$

As yet unconfirmed....



Oscillation physics:

MiniBooNE will check the LSND result with similar L/E,
higher statistics, and different systematics for the ν flux and particle ID

Non-oscillation physics:

- Charged current quasi-elastic
- Neutral current π^0
- Neutral current elastic scattering

BOONE COLLABORATION

University of Alabama
Bucknell University
University of Cincinnati
University of Colorado
Columbia University

Embry Riddle Aeronautical University
Fermi National Accelerator Laboratory

Indiana University
Los Alamos National Laboratory


Louisiana State University
University of Michigan
Princeton University

Y. Liu, I. Stancu
S. Koutsoliotas
E. Hawker, R.A. Johnson, **J.L. Raaf**
T. Hart, **R.H. Nelson**, E.D. Zimmerman
A.A. Aguilar-Arevalo, L. Bugel, L. Coney, J.M. Conrad,
J. Formaggio, J. Link, **J. Monroe**, **D. Schmitz**, M.H. Shaevitz,
M. Sorel, G.P. Zeller
D. Smith
L. Bartoszek, C. Bhat, S.J. Brice, B.C. Brown, D.A. Finley,
B.T. Fleming, R. Ford, F.G. Garcia, P. Kasper, T. Kobilarcik,
I. Kourbanis, A. Malensek, W. Marsh, P. Martin, F. Mills,
C. Moore, P. Nienaber, E. Prebys, A.D. Russell, P. Spentzouris,
R. Stefanski, T. Williams
D.C. Cox, J.A. Green, H. Meyer, R. Tayloe
G.T. Garvey, C. Green, W.C. Louis, G. McGregor, S. McKenney,
G.B. Mills, V. Sandberg, B. Sapp, R. Schirato, R. Van de Water,
N. Walbridge, D.H. White
R. Imlay, W. Metcalf, M. Sung, M. Wascko
J. Cao, Y. Liu, B.P. Roe, H.J. Yang
A.O. Bazarko, P.D. Meyers, **R.B. Patterson**,
F.C. Shoemaker, H.A. Tanaka

BOONE BEAM & DETECTOR

Beam

- 8 GeV protons from the Fermilab Booster directed into horn containing 71 cm Be target
- Secondary particles from target interactions (π , K) focused into 50 m decay region ($\pi \rightarrow \mu \nu_\mu$)
Absorber and 450 m of dirt "clean" the beam of everything except ν 's




Detector


- 12 m diameter sphere
- 8 inch Hamamatsu PMT's
- 1280 PMT's in signal region (10% coverage)
- 240 PMT's in veto region
- ~800 tons pure mineral oil

BEAM COMPOSITION


- protons on beryllium



- yield a high flux of ν_μ




- with a low ν_e background



ν_e background comparable to oscillation signal → need to know flux very well!!

- detailed simulations (GEANT3/GEANT4)
- CERN HARP measurements with MiniBooNE target replica
- BNL E910 production data with thin Be target
- Off-axis muon counter (LMC) → background ν_e 's from K decays
- 25/50 meter decay region → background ν_e 's from μ decays

DETECTOR CALIBRATION



Michel electrons (from cosmic μ decay)

Known energy spectrum between 0 and 52.3 MeV

Fix detector energy scale

Energy reconstruction accuracy: 14.8% at 52.3 MeV



4 Laser flasks

Known wavelength and intensity


Hit reconstruction:

PMT time/charge resolution

pre/after-pulsing

Oil optical properties:

attenuation, scattering,
surface reflections



7 Scintillator cubes

Optically isolated from oil

Detect stopping cosmic μ

Path length from muon hodoscope

Independent measurement of

muon energy up to 400 MeV


PARTICLE IDENTIFICATION

Veto Activity



Track Extent

Ring Profile




Hit Topology






no




possible




no



BEAM NEUTRINO EVENTS



- Beam sent to MiniBooNE in 1.6 μs wide spills
- DAQ triggers on FNAL Booster signal 4.6 μs before beam reaches target
- 19.2 μs window recorded surrounding each spill




So far....


204,000 ν_μ events from 1.8×10^{20}
protons on target

EARLY PHYSICS


Charged Current
Quasi-Elastic scattering
(CCQE)



Neutral Current
 π^0 production
(NC π^0)



Neutral Current
Elastic scattering
(NCE)




At MiniBooNE energies:

Charged current quasi-elastic: 39%

Neutral current π^0 production: 7%


Neutral current elastic: 17%



- ν_μ interactions only
- $\sim 10^{20}$ protons on target analyzed
- Data/MC comparisons relatively normalized
- Systematic uncertainties are preliminary

CHARGED CURRENT QUASI-ELASTIC SCATTERING

- ν_e CCQE events with similar kinematics are main signal for $\nu_\mu \rightarrow \nu_e$ oscillation search
- MiniBooNE is sensitive to ν_μ disappearance for $\Delta m^2 \sim 0.1\text{-}10 \text{ eV}^2$
reconstructible ν_μ energy and reasonably well-known cross section
- Characterize nuclear effects in ν -A interactions at $\sim 1 \text{ GeV}$



Event selection:

single muon-like $\bar{\nu}$ Cerenkov ring
scintillation light consistent with CCQE
high statistics: 30,000 events, 88% purity

Use QE kinematics to reconstruct E_ν from E_μ , $\cos \theta_\mu$
 $\sim 15\%$ energy resolution

Energy dependence sensitive to ν_μ disappearance

Data show reasonable agreement with MC predictions

CCQE: Q^2 DISTRIBUTION

Leptonic four-momentum transfer


$$Q^2 = -(p_\nu - p_\mu)^2$$

MC predictions based on Fermi gas
nuclear model (NUANCE)

Nuclear effects expected at low Q^2

Hint to more cross section suppression at low
 Q^2 in data compared to MC predictions

Physics or detector effect?



Low Q^2 suppression also seen in charged-current inclusive distributions in K2K near detectors, on O and Fe targets. (Ishida, NuInt01)

NEUTRAL CURRENT π^0 PRODUCTION

Background to $\nu_\mu \rightarrow \nu_e$ oscillation search


Knowledge of NC π^0 cross section crucial for distinguishing $\nu_\mu \rightarrow \nu_\tau$ from $\nu_\mu \rightarrow \nu_{\text{sterile}}$ in atmospheric neutrinos

Total cross section measurement and π^0 angular distribution constrain mechanisms for NC π^0 production


Production mechanisms:

Resonant: Nucleon goes into excited state (Δ, N) and decays by radiating π^0

Coherent: Neutrino scatters from entire nucleus. Nucleus remains in ground state and does not break apart.



NC π^0 : RECONSTRUCTION



FIT EVENT ASSUMING TWO RINGS (14 PARAMETERS)

- › decay vertex (4)
- › direction of γ 's (4)
- › mean emission points (2)
- › amount of Čerenkov/scintillation light (4)
- › no (e/ μ) ring ID

DETERMINE EVENT KINEMATICS (USING ČERENKOV LIGHT)

$$mc^2 = \sqrt{2 E_1 E_2 (1 - \cos \theta_{12})}$$

$$\vec{p} = E_1 \hat{u}_1 + E_2 \hat{u}_2$$

$$\beta \cos \theta_{CM} = \frac{|E_1 - E_2|}{E_1 + E_2}$$

NC π^0 : MASS DISTRIBUTION

Event selection from beam triggers:

$$N_{\text{TANK}} > 200$$

$$N_{\text{VETO}} < 6$$

$$R < 500 \text{ cm}$$


No decay electrons

$$E_{\gamma 1}, E_{\gamma 2} > 40 \text{ MeV}$$

Fitted curves MC-based parameterizations

Background peak near m_{π^0} expected

- final state interactions
- multi-pion events



Bin data in kinematic quantities

π^0 momentum (p_{π^0})

Energy asymmetry ($\beta \cos \theta_{CM} = \frac{|E_1 - E_2|}{E_1 + E_2}$)

Angle of π^0 relative to beam ($\cos \theta_{\pi^0}$)



Extract binned yields

NC π^0 : KINEMATIC DISTRIBUTIONS

π^0 momentum

Good data/MC agreement


Fall-off at high momentum due to
neutrino flux_v
overlapping \bar{C} rings



π^0 decay energy asymmetry



θ_{CM} - angle between π^0 decay axis in CM and π^0 direction in lab

$$\beta \cos \theta_{CM} = \frac{|E_1 - E_2|}{E_1 + E_2}$$



Fall-off due to γ energy cut

NC π^0 : KINEMATIC DISTRIBUTIONS



MC assumes Rein-Sehgal cross sections

Recent theories (Paschos, hep-ph/0309148) and experiments (K2K) suggest lower contribution from coherent pion production

MiniBooNE will extract coherent contribution

π^0 lab production angle

sensitive to production mechanism

coherent: forward-peaked

resonant: not as forward

NEUTRAL CURRENT ELASTIC SCATTERING


"Nucleon spin crisis"

What carries the proton spin? valence quarks, sea quarks, or gluons?

$\sigma(\text{NCE})/\sigma(\text{CCQE})$ ratio probes strange sea contribution to nucleon spin

Measure $\sigma(\text{NCE})$

Will help in understanding scintillation light for MiniBooNE oscillation search



$\nu_\mu + (p/n) \rightarrow \nu_\mu + (p/n)$

Typically sub- \bar{C} : dominated by scintillation



Low hit multiplicity, large scintillation fraction

Large cross section ($\sim 17\%$)

NEUTRAL CURRENT ELASTIC SCATTERING

Background subtraction:

- Beam excess clearly visible for < 150 tank hits
- Non-beam background due to decay electrons environmental activity
- Subtract with random triggers ("strobe trigger")



Event selection:

$$50 < N_{\text{TANK}} < 150 \text{ (50 hit threshold for vertex fit)}$$

$$\text{Scintillation light fraction} > 0.5$$

$$\text{Normalize MC to events with } N_{\text{TANK}} > 50$$

Reasonable agreement between data/MC for $N_{\text{TANK}} > 50$ with/without scintillation cut

ONGOING AND UPCOMING

Charged current quasi-elastic:

- compare with flux predictions
- ν_μ disappearance analysis
- probe low Q^2 region

Neutral current π^0 production:

- measure cross section
- analyze coherent contribution

Neutral current elastic:

- measure $\sigma_{\text{NC}}/\sigma_{\text{CC}}$ vs. Q^2
- probe Δs

MiniBooNE has been collecting data for > 1 year
 1.8×10^{20} protons on target
204K contained neutrino candidates

Detector working as expected

Reconstruction algorithms working well

STAY TUNED FOR MORE!